

About the Q^2 dependence of asymmetry A_1

A.V.Kotikov ¹

*Laboratoire de Physique Theorique ENSLAPP
LAPP, B.P. 110, F-74941, Annecy-le-Vieux Cedex, France*

and

D.V.Peshekhonov ²

Particle Physics Laboratory, JINR, Dubna, Russia.

Abstract

We analyse the proton and neutron data on spin dependent asymmetry $A_1(x, Q^2)$ supposing the DIS structure functions $g_1(x, Q^2)$ and $F_3(x, Q^2)$ have the similar Q^2 -dependence. As a result, we have obtained $\Gamma_1^p - \Gamma_1^n = 0.192$ at $Q^2 = 10 \text{ GeV}^2$ and $\Gamma_1^p - \Gamma_1^n = 0.165$ at $Q^2 = 3 \text{ GeV}^2$, in the best agreement with the Bjorken sum rule predictions.

PACS number(s): 13.60.Hb, 11.55.Hx, 13.88.+e

¹ On leave of absence from Particle Physics Laboratory, JINR, Dubna, Russia.
E-mail: kotikov@lapphp0.in2p3.fr; kotikov@sunse.jinr.dubna.su

² E-mail: dimitri@na47sun05.cern.ch; peshehon@sunse.jinr.dubna.su

An experimental study of the nucleon spin structure is realized by measuring of the asymmetry $A_1(x, Q^2) = g_1(x, Q^2)/F_1(x, Q^2)$. The most known theoretical predictions on spin dependent structure function $g_1(x, Q^2)$ of the nucleon were done by Bjorken [1] and Ellis and Jaffe [2] for the so called *first moment value* $\Gamma_1 = \int_0^1 g_1(x) dx$.

The calculation of the Γ_1 value requires the knowledge of structure function g_1 at the same Q^2 in the whole x range. Experimentally asymmetry A_1 is measuring at different values of Q^2 for different x bins. An accuracy of the past and modern experiments [3] - [9] allows to analyze data in the assumption [10]-[11] that asymmetry $A_1(x, Q^2)$ is Q^2 independent (structure functions g_1 and F_1 have the same Q^2 dependence). But the tune checking of the Bjorken and Ellis - Jaffe sum rules requires considering the Q^2 dependence of A_1 or g_1 (for recent studies of the Q^2 dependence of A_1 see [10]-[15]).

This article is based on our observation³ that the Q^2 dependence of g_1 and the spin average structure function F_3 is the same in a wide x range: $10^{-2} < x < 1$. At small x it seems that may be not true (see [16], [17]-[20]).

To demonstrate the validity of the observation, let's consider the nonsinglet (NS) Q^2 evolution of structure functions F_1 , g_1 and F_3 . The DGLAP equation for the NS part of these functions can be presented as⁴ :

$$\begin{aligned}\frac{dg_1^{NS}(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\gamma_{NS}^-(x, \alpha) \times g_1^{NS}(x, Q^2), \\ \frac{dF_1^{NS}(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\gamma_{NS}^+(x, \alpha) \times F_1^{NS}(x, Q^2), \\ \frac{dF_3^{NS}(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\gamma_{NS}^-(x, \alpha) \times F_3^{NS}(x, Q^2),\end{aligned}\tag{1}$$

where symbol \times means the Mellin convolution. Functions γ_{NS}^\pm are the reverse Mellin transforms of the anomalous dimensions $\gamma_{NS}^\pm(n, \alpha) = \alpha\gamma^{(0)}(n)_{NS} + \alpha^2\gamma_{NS}^{\pm(1)}(n) + O(\alpha^3)$ and the Wilson coefficients⁵ $\alpha b^\pm(n) + O(\alpha^2)$:

$$\gamma_{NS}^\pm(x, \alpha) = \alpha\gamma_{NS}^{(0)}(x) + \alpha^2\left(\gamma_{NS}^{\pm(1)}(x) + 2\beta_0 b^\pm(x)\right) + O(\alpha^3),\tag{2}$$

where $\beta(\alpha) = -\alpha^2\beta_0 - \alpha^3\beta_1 + O(\alpha^4)$ is QCD β -function.

The above mentioned Mellin transforms mean that

$$f(n, Q^2) = \int_0^1 dx x^{n-1} f(x, Q^2),\tag{3}$$

where $f = \{\gamma_{NS}^{(0)}, \gamma_{NS}^{\pm(1)}, b_{NS}^\pm, \gamma_{ij}^{(k)}, \gamma_{ij}^{*(k)}, b_i$ and $b_i^*\}$ with $k = 1, 2$ and $\{i, j\} = \{S, G\}$.

³The conclusion connects with our previous analysis [14].

⁴We use $\alpha(Q^2) = \alpha_s(Q^2)/4\pi$.

⁵Because we consider here the structure functions themselves but not the quark distributions. Note that more standard definition of $b_{NS}^+(n)$ and $b_{NS}^-(n)$ are $b_{1,NS}(n) = b_{2,NS}(n) - b_{L,NS}(n)$ and $b_{3,NS}(n)$.

Eqs. (1) show the Q^2 dependence of NS parts of g_1 and F_3 is the same (at least in first two orders of the perturbative QCD [21]) and differs from F_1 already in the first subleading order ($\gamma_{NS}^{+(1)} \neq \gamma_{NS}^{-(1)}$ [22] and $b_{NS}^+ - b_{NS}^- = (8/3)x(1-x)$). For the singlet parts of g_1 and F_1 evolution equations are :

$$\begin{aligned}\frac{dg_1^S(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2} \left[\gamma_{SS}^*(x, \alpha) \times g_1^S(x, Q^2) + \gamma_{SG}^*(x, \alpha) \times \Delta G(x, Q^2) \right], \\ \frac{dF_1^S(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2} \left[\gamma_{SS}(x, \alpha) \times F_1^S(x, Q^2) + \gamma_{SG}(x, \alpha) \times G(x, Q^2) \right],\end{aligned}\quad (4)$$

where

$$\begin{aligned}\gamma_{SS}(x, \alpha) &= \alpha \gamma_{SS}^{(0)}(x) + \alpha^2 \left(\gamma_{SS}^{(1)}(x) + b_G(x) \times \gamma_{GS}^{(0)}(x) + 2\beta_0 b_S(x) \right) \\ &\quad + O(\alpha^3), \\ \gamma_{SG}(x, \alpha) &= \frac{e}{f} \left[\alpha \gamma_{SG}^{(0)}(x) + \alpha^2 \left(\gamma_{SG}^{(1)}(x) + b_G(x) \times (\gamma_{GG}^{(0)}(x) - \gamma_{SS}^{(0)}(x)) + 2\beta_0 b_G(x) \right. \right. \\ &\quad \left. \left. + b_S(x) \times \gamma_{SG}^{(0)}(x) \right) \right] + O(\alpha^3)\end{aligned}$$

where $e = \sum_i^f e_i^2$ is the sum of charge squares of f active quarks. The equations for polarized anomalous dimensions $\gamma_{SS}^*(x, \alpha)$ and $\gamma_{SG}^*(x, \alpha)$ are similar. They can be obtained by replacing $\gamma_{Si}^{(1)}(x) \rightarrow \gamma_{Si}^{*(1)}(x)$ and $b_i(x) \rightarrow b_i^*(x)$ ($i = \{S, G\}$).

Note here the gluon term is not negligible for F_1 at $x < 0.1$ but for g_1 we can neglect the gluons for $x > 0.03$ [13]-[17]. The value $b_s^*(x)$ ($b_s(x)$) coincides with $b^-(x)$ ($b^+(x)$). The difference between $\gamma_{NS}^{-(1)}$ and $\gamma_{SS}^{*(1)} + b_G^*(x) \times \gamma_{GS}^{(0)}(x)$ is negligible due to its difference having no a power singularity at $x \rightarrow 0$ (i.e. no a singularity for them momentum transforms at $n \rightarrow 1$ in momentum space) and decreases as $O(1-x)$ at $x \rightarrow 1$ [26] (see also [27]). Contrary to this, the difference between $\gamma_{SS}^{(1)} + b_G(x) \times \gamma_{GS}^{(0)}(x)$ and $\gamma_{SS}^{*(1)} + b_G^*(x) \times \gamma_{GS}^{(0)}(x)$ contains the power singularity at $x \rightarrow 0$ (see [28, 21]).

This observation allows us to conclude the function :

$$A_1^*(x) = \frac{g_1(x, Q^2)}{F_3(x, Q^2)} \quad (5)$$

should be practically Q^2 independent at $x > 0.01$. Because the r.h.s. of Eqs.(1) and (4) contain integrals of structure functions, the approximate validity of (5) is supported also by the same x -dependence of $g_1(x, Q^2)$ and $F_3(x, Q^2)$ at fixed Q^2 . The asymmetry A_1 at $Q^2 = \langle Q^2 \rangle$ can be defined than as :

$$A_1(x_i, \langle Q^2 \rangle) = \frac{F_3(x_i, \langle Q^2 \rangle)}{F_3(x_i, Q_i^2)} \cdot \frac{F_1(x_i, Q_i^2)}{F_1(x_i, \langle Q^2 \rangle)} \cdot A_1(x_i, Q_i^2), \quad (6)$$

where x_i (Q_i^2) means an experimentally measured value of x (Q^2).

We use SMC and E143 proton and deuteron data for asymmetry $A_1(x, Q^2)$ [6] - [9]. To get $F_1(x, Q^2)$ we take NMC parametrization for $F_2(x, Q^2)$ [23] and SLAC parametrization for $R(x, Q^2)$ [24] ($F_1 \equiv F_2/2x[1 + R]$). To get the values of $F_3(x, Q^2)$ we parametrize the CCFR data [25] as a function of x and Q^2 (see Fig.1).

First, using eq.5, we recalculate the SMC [6, 7] and E143 [8, 9] measured asymmetry of the proton and deuteron at $Q^2 = 10 \text{ GeV}^2$ and 3 GeV^2 , which are average Q^2 of these experiments respectively (results are shown in Fig.2, 3) and get the value of $\int g_1(x)dx$ through the measured x ranges (see Table 1).

To obtain the first moment values $\Gamma_1^{p(d)}$ we have used an original estimations of SMC and E143 for unmeasured regions [6] - [9]. Results on the Γ_1 values are shown in the Table 1.

Table 1. The first moment value of g_1 of the proton and deuteron.

$x_{min} - x_{max}$	$< Q^2 >$	target type	$\int_{x_{min}}^{x_{max}} g_1 dx$	Γ_1	experiment
.003 – 0.7	10 GeV^2	proton	0.130	0.135	SMC
.003 – 0.7	10 GeV^2	deuteron	0.038	0.0362	SMC
.029 – 0.8	3 GeV^2	proton	0.123	0.130	E143
.029 – 0.8	3 GeV^2	deuteron	0.043	0.044	E143

As the last step we calculate the difference $\Gamma_1^p - \Gamma_1^n = 2\Gamma_1^p - (\Gamma_1^p + \Gamma_1^n)$ where $\Gamma_1^p + \Gamma_1^n = 2\Gamma_1^d/(1 - 1.5 \cdot \omega_D)$ and $\omega_D = 0.05$ [7, 9] .

At $Q^2 = 10 \text{ GeV}^2$ we get the following results :

$$\begin{aligned}
\Gamma_1^p - \Gamma_1^n &= 0.199 \pm 0.038 && (\text{SMC [7]}) \\
\Gamma_1^p - \Gamma_1^n &= 0.192 && (\text{our result}) \\
\Gamma_1^p - \Gamma_1^n &= 0.187 \pm 0.003 && (\text{Theory})
\end{aligned} \tag{7}$$

and at $Q^2 = 3 \text{ GeV}^2$:

$$\begin{aligned}
\Gamma_1^p - \Gamma_1^n &= 0.163 \pm 0.026 && (\text{E143 [9]}) \\
\Gamma_1^p - \Gamma_1^n &= 0.165 && (\text{our result}) \\
\Gamma_1^p - \Gamma_1^n &= 0.171 \pm 0.008 && (\text{Theory})
\end{aligned} \tag{8}$$

As a conclusion, we would like to note

- our observation that function $A_1^*(x)$ is Q^2 independent at large and intermediate x is supported by good agreement (see Fig. 2,3) of present analysis with other estimations [12]-[16] of the Q^2 dependence of the A_1 ;
- at small x structure functions $g_1(x, Q^2)$ and $F_3(x, Q^2)$ may have the same behaviour too (in traditional, Regge-motivated consideration $f \sim x^\delta$, where $\delta \geq 0$) [30] ⁶.

⁶According to the recent analysis [17]-[20], however, the situation may be more complicated.

- The value of $\Gamma_1^p - \Gamma_1^n$ obtained in the supposition that g_1 and F_3 have the same Q^2 dependence improves the agreement with the Bjorken sum rule prediction.

Acknowledgements

We are grateful to W.G. Seligman for providing us the available CCFR data of Ref.[25] and A.V. Efremov for discussions.

This work is supported partially by the Russian Fund for Fundamental Research, Grant N 95-02-04314-a.

References

- [1] J. D. Bjorken, Phys. Rev., **148** (1966) 1467; **D1** (1970) 1376.
- [2] J. Ellis and R. L. Jaffe, Phys. Rev., **D9** (1974) 1444; **D10** (1974) 1669.
- [3] SLAC-E80 Collaboration, M. J. Alguard *et al.*, Phys. Rev. Lett. **37** (1976) 1261; **41** (1978) 70;
SLAC-E130 Collaboration, G. Banm *et al.*, Phys. Rev. Lett. **51** (1983) 1135.
- [4] EM Collaboration, J. Ashman *et al.*, Phys. Lett. **B206** (1988) 364; Nucl. Phys. **B328** (1989) 1.
- [5] SLAC-E142 Collaboration, P. L. Anthony *et al.*, Phys. Rev. Lett. **71** (1993) 959.
- [6] SM Collaboration, B. Adams *et al.*, Phys. Lett. **B329** (1994) 399.
- [7] SM Collaboration, B. Adams *et al.*, Phys. Lett. **B357** (1995) 248.
- [8] SLAC-E143 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **74** (1995) 346.
- [9] SLAC-E143 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **75** (1995) 25.
- [10] J. Ellis and M. Karliner, Phys. Lett. **B313** (1993) 131.
- [11] F. E. Close, Report No RAL-93-034, 1994;
F. E. Close and R. G. Roberts, Phys. Lett. **B316** (1993) 165.
- [12] G. Altarelli, P. Nason, and G. Ridolfi, Phys. Lett. **B320** (1994) 152;
- [13] T. Gehrmann and W. J. Stirling, Z. Phys. **C65** (1995) 470.
- [14] A. V. Kotikov and D. V. Peshekhonov, Reports ENSLAPP-A-499/94, 1994;
ENSLAPP-A-59/96, 1996.
- [15] SLAC-E143 Collaboration, K. Abe *et al.*, Phys. Lett. **B364** (1995) 61.

- [16] M. Gluck *et al*, Phys. Lett. **B359** (1995) 210, Report No. DO-Th 95/13, 1995 (hep-ph/9508347).
- [17] R. D. Ball, S. Forte, and G. Ridolfi, Nucl. Phys. **B444** (1995) 287 (E: **B449** (1995) 680); Report CERN-Th/95-266, Edinburg 95/556, GeF-TH-9/95 (hep-ph/9510449);
T. Gehrmann and W. J. Stirling, Report No DTP/95/62.
- [18] S. J. Brodsky, M. Burkardt, and I. Schmidt, Nucl. Phys. **B441** (1995) 197.
- [19] S. Forte, Report No CERN-Th/95-305 (hep-ph/9511345);
R. D. Ball, Report No Edinburg 95/558 (hep-ph/9511330);
T. Morii, S. Tanaka, and T. Yamanishi, Report No KOBE-FHD-95-05 (hep-ph/9508231);
T. Gehrmann and W. J. Stirling, Reports No DTP/95/78 (hep-ph/9510243) and DTP/95/82 (hep-ph/9512406).
- [20] B. I. Ermolaev, S. I. Manayenkov, and M. G. Ryskin, Z. Phys. **C69** (1996) 259;
J. Bartels, B. I. Ermolaev, and M. G. Ryskin, Reports DESY-95-124, DESY-96-025;
J. Blumlein and A. Vogt, Phys. Lett. **B370** (1996) 149; Reports DESY-96-050, DESY-96-041.
- [21] J. Kodaira *et al.*, Phys. Rev. D **20** (1979) 627; Nucl. Phys. **B159** (1979) 99.
- [22] D. A. Ross and C. T. Sachrajda, Nucl. Phys. **B149** (1979) 497;
G. Altarelli, Phys. Rep. **81** (1982) 1.
- [23] NM Collaboration, P. Amandruz *et al.*, Phys. Lett. **B295** (1992) 159.
- [24] SLAC Collaboration, L. W. Whitlow *et al.*, Phys. Lett. **B250** (1990) 193.
- [25] CCFR Collab. P.Z. Quintas *et al*, *Phys. Rev. Lett.* **71** (1993) 1307;
CCFR Collab. M. Shaevitz *et al.* *Nucl. Phys. Proc. Suppl.* **B38** (1995) 188;
P.Z. Leung, Columbia Univ. Thesis R-1478, CU-382, Nevis-276 (1992);
W.C. Quintas, Columbia Univ. Thesis R-1482, CU-383, Nevis-277 (1992).
- [26] R. Merting and W. L. van Neervan, Report No INLO-PUB-6/95, NIKHEF-H/95-31 (hep-ph/9506451).
- [27] W. Vogelsang, Repors RAL-TR-95-071 (hep-ph/9512218); RAL-TR-96-020 (hep-ph/9603366).
- [28] A. Gonzalez-Arroyo, C. Lopez, and F. J. Yndurain, Nucl. Phys. **B174** (1980) 474.
- [29] C. Bourrely and J. Soffer, Nucl. Phys. **B445** (1995) 341.
- [30] R. L. Heimann, Nucl. Phys. **B64** (1973) 429;
J. Ellis and M. Karliner, Phys. Lett. **B213** (1988) 73.

Figure Captions

Figure 1. DIS structure function $F_3(x, Q^2)$. CCFR data [25] and the parametrization.

Figure 2. SMC [6] and E143 [8] measured virtual photon-nucleon asymmetry $A_1^p(x)$ as a function of x (shown as a close points) in comparison with evolved to $Q^2 = 10$ (3) GeV^2 , respectively (open points).

Figure 3. SMC [7] and E143 [9] measured virtual photon-nucleon asymmetry $A_1^d(x)$ as a function of x (shown as a close points) in comparison with evolved to $Q^2 = 10$ (3) GeV^2 , respectively (open points).





